

A Bicategory Approach to Differential Cohomology

Markus Upmeyer

December 27, 2012

Abstract

A very natural bicategory approach to differential cohomology is presented. Based on the axioms of Bunke-Schick, a symmetric monoidal groupoid is associated to any differential cohomology theory. The main result is then that such a differential refinement is unique up to equivalence of the corresponding symmetric monoidal groupoids. The uniqueness results for rationally-even theories are interpreted in this framework. Moreover, we show how the bicategory formalism may be used to give a simple construction of a differential refinement for any generalized cohomology theory.

Acknowledgements. I would particularly like to thank Ralf Meyer for many fruitful discussions out of which this paper grew.

1 Introduction

Differential cohomology is a refinement of generalized cohomology that takes gauge-theoretic features of the underlying manifold into account. Let $(\hat{E}^*, R, I, a)$ denote a differential extension of a generalized cohomology theory E^* in the sense of [BS10]. Recall that this means the following:

Suppose E is a cohomology theory and let $V := E^*(pt) \otimes \mathbb{R}$ (a graded vector space). Let **Manifolds** denote the category of smooth manifolds. A *differential extension* $\hat{E}$ of E is a family of contravariant functors

$$\hat{E}^* : \mathbf{Manifolds} \longrightarrow \mathbf{Abel}, \quad (* \in \mathbb{Z})$$

into the category of Abelian groups together with natural transformations

$$\begin{aligned}\Omega^{*-1}(-; V)/\text{im}(d) &\xrightarrow{a} \hat{E}^* \\ \hat{E}^* &\xrightarrow{R} \Omega^*(-; V) \\ \hat{E}^* &\xrightarrow{I} E^*(-)\end{aligned}$$

such that the following diagram commutes for any $M \in \text{Ob}(\mathbf{Manifolds})$ and has an exact upper horizontal line

$$\begin{array}{ccccccc} E^{*-1}(M) & \xrightarrow{ch} & \Omega^{*-1}(M; V)/\text{im}(d) & \xrightarrow{a} & \hat{E}^*(M) & \xrightarrow{I} & E^*(M) \longrightarrow 0 \\ & & \searrow d & & \downarrow R & & \downarrow ch \\ & & & & \Omega_{cl}^*(M; V) & \longrightarrow & H_{sing}^*(M; V) \end{array}$$

Here we have used differential forms and singular cohomology with coefficients in a graded vector space. Of course, an *isomorphism of differential cohomology theories* is defined to be a natural transformation which is compatible with the maps a, R, I . One of the main results in [BS10] is the following:

Theorem 1 ([BS10]). *Suppose E is rationally even, i.e. E is a cohomology theory whose graded coefficient group $E^*(pt)$ is torsion in all odd degree. If $E^*(pt)$ is countably generated in all degrees then any two differential extensions $\hat{E}$ with integration will be isomorphic via a unique isomorphism respecting integration.*

2 Uniqueness of Differential Extensions

For every manifold M , the transformation $a : \Omega^{*-1}(M; V)_d \longrightarrow E^*(M)$ may be used to define a groupoid

$$\tilde{E}^*(M) = \begin{cases} \text{objects} & x \in E^*(M), \\ \text{morphisms from} & \eta \in \Omega^{*-1}(M; V)_d \text{ with } a(\eta) = y - x. \\ x \in E^*(M) \text{ to } y \in E^*(M) \end{cases}$$

Here, $\Omega^{*-1}(M; V)_d$ denotes the quotient of $\Omega^{*-1}(M; V)$ by $\text{im}(d)$. Using addition, $\tilde{E}^*(M)$ becomes a strict symmetric monoidal groupoid. For varying

M these may be organized into a functor

$$\tilde{E}^* : \mathbf{Manifolds} \longrightarrow \mathbf{SymMonCat}_{\text{str}}.$$

Similarly, $d : \Omega^{*-1}(M; V)_d \longrightarrow \Omega^*(M; V)$ and $\delta : C'^{-1}(M; V)_\delta \rightarrow C^*(M; V)$ may be used to construct functors

$$\tilde{\Omega}^*, \mathbb{C}_{\text{sing}}^* : \mathbf{Manifolds} \longrightarrow \mathbf{SymMonCat}_{\text{str}}.$$

The transformation R along with the property $R \circ a = d$ yields a natural transformation

$$R : \tilde{E}^* \rightarrow \tilde{\Omega}^*.$$

Proposition 2. *Let $\hat{E}^*$ and $\hat{E}'^*$ be differential extensions of E^* . For each $M \in \mathbf{Manifolds}$, there exists an equivalence of the corresponding symmetric monoidal groupoids*

$$\Phi_M : \tilde{E}^*(M) \longrightarrow \tilde{E}'^*(M)$$

which is natural in M .

Proof. Represent E^* by an Ω -spectrum (E_n, ε_n) . Fix a degree n . By Proposition 2.3. in [BS10] there exists a sequence of pointed manifolds $\mathcal{E}_n^{(i)}$, $i \in \mathbb{N}$, together with pointed embeddings of submanifolds $\kappa_i : \mathcal{E}_n^{(i)} \hookrightarrow \mathcal{E}_n^{(i+1)}$ as well as $(i-1)$ -connected pointed maps $x_i : \mathcal{E}_n^{(i)} \rightarrow E_n$ such that $x_{i+1} \circ \kappa_i = x_i$ and such that for any finite dimensional CW-Complex $X \in \mathbf{CW}^+$ the canonical map

$$\text{colim}_i [X, \mathcal{E}_n^{(i)}] \rightarrow [X, E_n] = E^n(X)$$

is an isomorphism.

One may then choose $\hat{u}_i \in \hat{E}^n(\mathcal{E}_n^{(i)})$ with

$$\kappa_i^* \hat{u}_{i+1} = \hat{u}_i, \quad I(\hat{u}_i) = [x_i], \quad \text{Rham}(R(\hat{u}_i)) = ch[x_i]$$

Similarly, one may choose $\hat{u}'_i \in \hat{E}'^n(\mathcal{E}_n^{(i)})$. We are now ready to define Φ_M on objects: For $\hat{v} \in \hat{E}^n(M)$ write $I(\hat{v}) = [M \xrightarrow{f} E_n]$ for some f and factor $f \simeq x_i \circ f_i$ up to homotopy for some $f_i : M \rightarrow \mathcal{E}_n^{(i)}$. Then

$$\hat{v} - f_i^* \hat{u}_i = a(\alpha) \text{ for some } \alpha \in \Omega^{n-1}(M; V)_d$$

and we define

$$\Phi_M(\hat{v}) := f_i^*(\hat{u}'_i) + a'(\alpha).$$

As is shown in [BS10], Φ_M is well-defined, compatible with I, R, a and natural in M . We use the identity on $\Omega^{n-1}(M; V)_d$ to view Φ_M as a functor

$$\Phi_M : \tilde{E}^n(M) \rightarrow \tilde{E}^n(M),$$

which is clearly full and faithful. It is also essentially surjective: For $\hat{u}' \in \hat{E}'^*(M)$ choose $\hat{u} \in \hat{E}^*(M)$ with $I(\hat{u}) = I'(\hat{u}')$. Then $\Phi_M(\hat{u}) - \hat{u}' = a'(\alpha)$ for some $\alpha \in \Omega^{n-1}(M; V)_d$ and then $\hat{u} - a(\alpha)$ maps to $\hat{u}'$.

Thus Φ_M is an equivalence of categories. It will therefore suffice to show that Φ_M is a symmetric monoidal functor. The deviation of Φ_M from being additive is a natural transformation

$$(\Phi_M \circ +) - (+' \circ (\Phi_M \times \Phi_M)) : \hat{E}^n(M) \times \hat{E}^n(M) \rightarrow \hat{E}^n(M)$$

that composes with I to zero, so that it factors over a natural transformation

$$B : \hat{E}^n(M) \times \hat{E}^n(M) \rightarrow \Omega^{n-1}(M; V)_d / \text{im}(ch).$$

satisfying

$$\begin{aligned} B(\hat{u}, \hat{v} + \hat{w}) + B(\hat{v}, \hat{w}) &= B(\hat{u}, \hat{v}) + B(\hat{u} + \hat{v}, \hat{w}), \\ B(\hat{u}, \hat{v}) &= B(\hat{v}, \hat{u}), \\ B(\hat{u}, 0) &= 0 = B(0, \hat{u}). \end{aligned}$$

Together with the identity arrows $0 \xrightarrow{0} \Phi_M(0) = 0$ the transformation B gives Φ_M the structure of a symmetric monoidal functor. The naturality of B – not in M but in the objects of $\hat{E}^n(M)$ follows from the observation that

$$\Phi_M(\hat{v} + a(\beta)) = \Phi_M(\hat{v}) + a(\beta).$$

□

Remark 3. Under additional assumptions (rationally even, finiteness conditions on the coefficients) the results of [BS10] amount to the assertion that one may even pick a *strict* monoidal equivalence.

3 Construction of Differential Extensions

3.1 The Refined Chern Character

Let E_n be an Ω -spectrum with structure maps $\varepsilon_n : \Sigma E_n \rightarrow E_{n+1}$ which are adjoint to homeomorphisms.

The fundamental groupoid $\text{Map}(X, E_n) = \Pi_1 E_n^X$ has a monoidal structure: there are two canonical addition maps

$$E_n \times E_n \approx \Omega^2 E_{n-2} \times \Omega^2 E_{n-2} \longrightarrow \Omega^2 E_{n-2} \approx E_n$$

given by “horizontal” and “vertical” concatenation which equip $\text{Map}(X, E_n)$ with two weak monoidal structures $\ominus, \oplus$. The associators and unitors are given by the usual homotopies. The identity object is given in both cases by the constant map $I : X \rightarrow E_n$ onto the basepoint. We have an *equality*

$$(f_1 \ominus g_1) \oplus (f_2 \ominus g_2) = (f_1 \oplus f_2) \ominus (g_1 \oplus g_2)$$

which together with the unitors yield a natural isomorphism $\ominus \Rightarrow \oplus$. Also, by an Eckmann-Hilton type argument we obtain a braid $\gamma_{f,g} : f \ominus g \Rightarrow g \ominus f$ that turns $(\text{Map}(X, E_n), \ominus, I, \gamma)$ into a symmetric monoidal category.

The deRham homomorphisms may be interpreted as a strict symmetric monoidal equivalence of categories

$$\rho_X : \tilde{\Omega}^n(M) \xrightarrow{\sim} \mathcal{C}_{sing}^n(M)$$

which are the components of a natural transformation $\tilde{\Omega}^n \rightarrow \mathcal{C}_{sing}^n$. Every component is an equivalence and we would like to invert ρ . To pick inverse equivalences systematically we use bicategories:

With monoidal transformations as 2-arrows **SymMonCat** forms a strict 2-category. We will view **Manifolds** as a 2-category with only identity 2-arrows. Then ρ is a strict transformation between the two strict functors $\tilde{\Omega}^n, \mathcal{C}_{sing}^n$ of bicategories.

Lemma 4. *Suppose $\mathcal{C}$ is a category, considered trivially as a bicategory. Let*

$$F, G : \mathcal{C} \longrightarrow \mathbf{SymMonCat}$$

be strict functors of bicategories and $u : F \Rightarrow G$ a strict transformation which has the property that all of the functors $u_X : FX \rightarrow GX$ are equivalences. Then there exists a weak transformation $v : G \Rightarrow F$ such that uv and vu are the identity functors up to modifications.

Proof. The monoidal functor u_X comes equipped with a natural transformation $\mu_X : \otimes(u_X \times u_X) \Rightarrow u_X \circ \otimes$. For any $X \in \mathcal{C}$ pick $v_X : GX \rightarrow FX$, $\varepsilon_X : 1 \Rightarrow u_X \circ v_X$, and $\eta_X : v_X \circ u_X \Rightarrow 1$ such that $(u_X, v_X, \varepsilon_X, \eta_X)$ form an adjoint equivalence. Using the unit η_X and counit ε_X of the adjunction we may make v_X symmetric weak monoidal: define the inverse of ν_X as

$$v_X \circ \otimes \xRightarrow{v(\varepsilon \otimes \varepsilon)} v_X \circ \otimes \circ (u_X v_X \times u_X v_X) \xRightarrow{v\mu(v \times v)} v_X \circ u_X \circ \otimes \circ (v_X \times v_X) \xRightarrow{\eta(v \times v)} \otimes \circ (v_X \times v_X)$$

The unit η_X and counit ε_X then become monoidal transformations. It remains to check that the monoidal functors (v_X, μ_X) depend functorially on X up to coherent 2-cells $v_f : (Ff, \phi) \circ (v_X, \nu_X) \Rightarrow (v_Y, \nu_Y) \circ (Gf, \gamma)$ for $f : X \rightarrow Y$ in $\mathcal{C}$. The monoidal transformation v_f is defined as the 2-cell

$$\begin{array}{ccccc} GX & \xrightarrow{1} & GX & \xrightarrow{Gf} & GY \\ & \searrow v_X & \downarrow \varepsilon_X & \nearrow u_Y & \downarrow \eta_Y \\ & & FX & \xrightarrow{Ff} & FY \\ & & & \nearrow u_X & \downarrow 1 \\ & & & & FY \end{array}$$

Note that the middle diagram commutes. The coherence conditions follow from the zig-zag equations for ε_X, η_X and are left to the reader. $\square$

We may thus find a *weak* transformation, unique up to modifications,

$$\tau : \mathcal{C}_{sing}^n \Longrightarrow \tilde{\Omega}^n.$$

Proposition 5. *It is possible to lift the Chern character transformation $ch : [M, E_n] \rightarrow H^n(M; V)$ to a weak monoidal functor*

$$\text{Map}(M, E_n) \rightarrow \mathcal{C}_{sing}^n(M)$$

natural in M , i.e. to a strict transformation $\text{Map}(-, E_n) \rightarrow \mathcal{C}_{sing}^n$.

Proof. Pick a compatible family $\iota_n \in C_{sing}^n(E_n; V)$ of fundamental cocycles implementing the Chern character: $\int_I \varepsilon_{n-1}^* \iota_n = \iota_{n-1}$. Define the functor on

objects by $f \mapsto f^* \iota_n$ and on morphisms by $[H] \mapsto \int_I H^* \iota_n$. The latter is welldefined since the morphisms in $\mathcal{C}_{sing}^n$ are $\text{im}(\delta)$ -cosets. Different choices of fundamental cocycles yield modifications of this transformation. The proof that this functor is monoidal is deferred to the appendix. $\square$

Composing this transformation with the weak inverse τ of the deRham transformation, we obtain a weak transformation

$$ch : \text{Map}(-, E_n) \Longrightarrow \tilde{\Omega}^n.$$

More explicitly, we are given symmetric monoidal functors

$$ch_M : \text{Map}(M, E_n) \longrightarrow \tilde{\Omega}^n(M),$$

i.e. a functor ch_M together with a natural transformation, denoted $ch_M^+ : ch_M \circ \Theta \rightarrow + \circ (ch_M \times ch_M)$, and a map $ch_M(\text{const}) \rightarrow 0$ which may be chosen to be zero. Also, for every smooth map $f : M \rightarrow N$ we have a map

$$ch_f : E_n^N \longrightarrow \Omega^{n-1}(M; V)_d$$

with

$$d(ch_f(c)) = f^* ch_N(c) - ch_M(c \circ f) \quad \text{for all } c : N \rightarrow E_n \quad (1)$$

$$f^* ch_N(H) + ch_f(H_0) = ch_f(H_1) + ch_M(H \circ (f \times id_I)) \quad \text{for a homotopy } H : H_0 \simeq H_1 \quad (2)$$

where $H : N \times I \rightarrow E_n$. The coherence conditions for $M \xrightarrow{f} N \xrightarrow{g} O \xrightarrow{c} E_n$ amount to

$$ch_{gf}(c) = ch_f(c \circ g) + f^* ch_g(c), \quad (3)$$

$$ch_{id} = 0. \quad (4)$$

The compatibility condition for the ι_n ensure that for a map $h : M \rightarrow E_{n-1}$ viewed as a homotopy $H : M \times I \rightarrow E_n$ from const to const we have

$$ch_M(h) = ch_M(H) \in \Omega^{n-1}(M; V)_d.$$

4 Construction of Differential Refinements

Consider the equivalence relation $\sim$ on the set $\mathcal{E}^n(M) = E_n^M \times \Omega^{n-1}(M; V)_d$ given by $(c_1, \omega_1) \sim (c_2, \omega_2)$ iff there exists a homotopy $H : c_1 \rightarrow c_2$ with

$$ch_M(H) = \omega_1 - \omega_2$$

Reflexivity, symmetry, and transitivity of $\sim$ follow from the functoriality properties of ch_M , respectively: that it preserves the identity, inverses of isomorphisms, and composition. We may now define

$$\hat{E}^n(M) = \mathcal{E}^n(M) / \sim.$$

For a smooth map $M \xrightarrow{f} N$ we let

$$\hat{E}^n(f) : \hat{E}^n(N) \longrightarrow \hat{E}^n(M), \quad [c, \omega] \mapsto [c \circ f, f^* \omega + ch_f(c)]$$

This is well-defined by (2) and yields a functor $\hat{E}^n : \mathbf{Manifolds} \longrightarrow \mathbf{Set}$ by (3) and (4). The group structure is given by

$$\begin{aligned} [c_1, \omega_1] + [c_2, \omega_2] &= [c_1 \ominus c_2, \omega_1 + \omega_2 + ch_M^+(c_1, c_2)], \\ 0 &= [\text{const}, 0]. \end{aligned}$$

Define natural transformations

$$\begin{aligned} \Omega^{n-1}(M; V)_d &\xrightarrow{a} \hat{E}^n(M), & a(\omega) &= [\text{const}, \omega] \\ \hat{E}^n(M) &\xrightarrow{I} E^n(M), & I[c, \omega] &= [c] \in [M, E_n] = E^n(M) \\ \hat{E}^n(M) &\xrightarrow{R} \Omega^n(M; V)_{d=0}, & R[c, \omega] &= ch(c) + d\omega \end{aligned}$$

The map R is well-defined by functoriality of ch_M .

Proposition 6. *$\hat{E}^n$ is a differential cohomology theory in the sense of [BS10]. That is, we have a commutative diagram with an exact row:*

$$\begin{array}{ccccccc} E^{n-1} & \xrightarrow{ch} & \Omega^{n-1}(-; V)_d & \xrightarrow{a} & \hat{E}^n & \xrightarrow{I} & E^n \longrightarrow 0 \\ & & \searrow d & & \downarrow R & & \downarrow ch \\ & & & & \Omega^n(-; V)_{d=0} & \longrightarrow & H_{dR}^n(-; V) \end{array}$$

Proof. The commutativity is clear. Now, if $I[c, \omega] = 0$ there exists a homotopy $H : c \rightarrow \text{const}$ so that $[c, \omega] = [\text{const}, \omega + ch_M(H)] = a(\omega + ch_M(H))$. Also $I \circ a = 0$. If $a(\omega) = [\text{const}, 0]$ then there exists $H : \text{const} \rightarrow \text{const}$ with $ch_M(H) = \omega$. and by construction $ch_M(H)$ represents $ch(H)$. Conversely, a map $c : M \rightarrow E_{n-1}$ may be viewed as a homotopy $H : \text{const} \simeq \text{const} : M \rightarrow E_n$. Then $a(ch(h)) = [\text{const}, ch(h)] = [\text{const}, 0]$, using H and the compatibility of the ι_n . $\square$

5 Appendix


The goal in this section is to show that for a fixed $X \in \mathbf{Manifolds}$

$$ch : \text{Map}(X, E_n) \longrightarrow \mathcal{C}_{sing}^n(X)$$

is a monoidal functor. Using $E_n \approx \Omega E_{n-1}$, the category $\text{Map}(X, E_n)$ is isomorphic to the category $\text{Map}(X \times (I, \partial I), (E_{n-1}, *))$ which carries an obvious weak monoidal structure. It will be of technical advantage to work instead with the isomorphic weak monoidal category $\mathcal{E}$ defined as follows:

In the sequel we will make reference to the face and degeneracy maps of the simplicial set $E_{n-1}^{X \times \Delta^\bullet}$.


Objects are $Ob(\mathcal{E}) = (E_{n-1}, *)^{X \times (\Delta^1, \partial \Delta^1)}$, typical elements of which are denoted by $f, g, h, \dots$. Morphisms $f \rightarrow g$ are homotopy classes of maps $H : X \times \Delta^2 \longrightarrow E_{n-1}$ with $d_2 H = f, d_1 H = g, d_0 H = *$. Here, a homotopy between arrows H_0 and H_1 from f to g is defined as a map $G : X \times \Delta^3 \longrightarrow E_{n-1}$ with $d_0 G = H_1, d_1 G = H_0, d_2 G = s_1 g, d_3 G = s_1 f$.



Just as for simplicial homotopy groups, composition is given by picking arbitrary horn fillers. The identity is $id_f = [s_1 f]$.


The monoidal structure is given by the fact that there are *canonical* horn fillers coming from canonical retractions $r_i : \Delta^n \rightarrow \Lambda_i^n$, $n \geq 2$ defined by assigning to $x \in \Delta^n$ the unique intersection point of Λ_i^n with the one-dimensional

affine subspace stationed at x in direction of the orthogonal complement of $d_i(\Delta^{n-1})$. We will only need $n = 2, 3$:




Now, given two objects $f, g \in \mathcal{E}$ we may glue them together to $X \times \Lambda_1^2 \rightarrow E_{n-1}$ and use $id_X \times r_2$ to canonically extend to $\sigma(f, g) : X \times \Delta^2 \rightarrow E_{n-1}$. Then set

$$f \oplus g = d_1 \sigma(f, g)$$




The tensor product of morphisms may be defined similarly by canonical extensions. Given $H_0 : f_0 \rightarrow g_0$ and $H_1 : f_1 \rightarrow g_1$ define $H_0 \oplus H_1 : f_0 \oplus f_1 \rightarrow g_0 \oplus g_1$ as follows:




Canonically fill the horn Λ_0^3 given by $(\mathbf{0}, \mathbf{2}, \mathbf{3}) = s_0 f_0$, $(\mathbf{2}, \mathbf{3}, \mathbf{4}) = s_0(f_0 \oplus f_1)$, $(\mathbf{0}, \mathbf{3}, \mathbf{4}) = \sigma(f_0, f_1)$ to obtain a 3-simplex $\tau = (\mathbf{0}, \mathbf{2}, \mathbf{3}, \mathbf{4})$. Next, canonically fill the horn Λ_1^3 given by $(\mathbf{0}, \mathbf{2}, \mathbf{4}) = d_0 \tau$, $(\mathbf{0}, \mathbf{1}, \mathbf{2}) = H_0$, $(\mathbf{0}, \mathbf{1}, \mathbf{4}) = s_0 f_1$ to a 3-simplex $\sigma = (\mathbf{0}, \mathbf{1}, \mathbf{2}, \mathbf{4})$. Finally, fill Λ_1^3 given by $(\mathbf{1}, \mathbf{2}, \mathbf{4}) = d_1 \sigma$, $(\mathbf{1}, \mathbf{2}, \mathbf{5}) = \sigma(g_0, g_1)$, $(\mathbf{1}, \mathbf{4}, \mathbf{5}) = H_1$ to a 3-simplex κ . Then $H_0 \oplus H_1 = d_1 \kappa$.

The identity object is $\ast = \text{const} : X \times (\Delta^1 \times \partial\Delta^1) \rightarrow (E_{n-1}, \ast)$. The left unitor is $\lambda_f^{-1} = \sigma(f, \ast) : f \rightarrow f \oplus \ast$. The right unitor is given as the side of an appropriately filled horn Λ_1^3 given by $(\mathbf{0}, \mathbf{1}, \mathbf{2}) = \sigma(\ast, f)$, $(\mathbf{1}, \mathbf{2}, \mathbf{3}) = s_1 f$, $(\mathbf{0}, \mathbf{1}, \mathbf{3}) = s_0 f$ as in the next figure. Let $\tau = (\mathbf{0}, \mathbf{1}, \mathbf{2}, \mathbf{3})$ denote the canonical filler. Then the right unitor is $\rho_f = d_1 \tau : \ast \oplus f \rightarrow f$.



The associator is defined as follows:



$(\mathbf{0}, \mathbf{1}, \mathbf{3}) = \sigma(g, f)$, $(\mathbf{0}, \mathbf{1}, \mathbf{4}) = \sigma(h \oplus g, f)$, $(\mathbf{0}, \mathbf{3}, \mathbf{4}) = \sigma(h, g)$ may be filled to a 3-simplex $\tau = (\mathbf{0}, \mathbf{1}, \mathbf{3}, \mathbf{4})$. Next, $(\mathbf{1}, \mathbf{3}, \mathbf{4}) = d_2\tau$, $(\mathbf{1}, \mathbf{2}, \mathbf{3}) = \sigma(h, g \oplus f)$, and $(\mathbf{2}, \mathbf{3}, \mathbf{4}) = s_0h$ may be filled to another 3-simplex $\kappa = (\mathbf{1}, \mathbf{2}, \mathbf{3}, \mathbf{4})$. Then


$$\alpha_{h,g,f} = d_2\kappa : (h \oplus g) \oplus f \longrightarrow h \oplus (g \oplus f)$$

Once we show that we have an isomorphism of categories $\text{Map}(X \times (I, \partial I), (E_{n-1}, *)) \cong \mathcal{E}$ that maps the tensor product functor, the unitors, and the associators onto each other it will be clear that $\mathcal{E}$ in fact is a monoidal category. The isomorphism is given by the identity on objects and by assigning to an arrow $H : f \rightarrow g$ in $\text{Map}(X \times (I, \partial I), (E_{n-1}, *))$ the arrow

$$h : X \times \Delta^2 \longrightarrow E_{n-1}, (s, t) \mapsto \begin{cases} H(s, t/s) & (s > 0), \\ * & (s = 0). \end{cases}$$

where here we view Δ^2 as $\{(s, t) \in I^2 \mid \frac{1-s}{2} \leq t \leq \frac{1+s}{2}\} \subset I^2$. The functor in the other direction is given on morphisms by a “vertical” retraction $I^2 \twoheadrightarrow \Delta^2$ mapping $I^2 \setminus \Delta^2$ onto $\partial\Delta^2$.

The following are a comparison of the two right unitors whose homotopy classes are clearly mapped onto each other.



References

- [BS10] U. Bunke and T. Schick, *Uniqueness of smooth extensions of generalized cohomology theories*, Journal of topology **3** (2010), 110–156.
- [CS85] J. Cheeger and J. Simons, *Differential characters and geometric invariants*, Geometry and topology (College Park), Lecture Notes in Mathematics, vol. 1167, Springer, Berlin, 1985, pp. 50–80.

- [HS05] M.J. Hopkins and I.M. Singer, *Quadratic functions in geometry, topology and M-theory*, J. Differential Geom. **70** (2005), 329–452.
- [SS08] J. Simons and D. Sullivan, *Axiomatic characterization of ordinary differential cohomology*, Journal of Topology **1** (2008), 45–56.
- [Upm] M. Upmeyer, *Differential cohomology, PhD-thesis in preperation, Georg-August University, Göttingen*.
- [Upm11] ———, *Products in Generalized Differential Cohomology*, arXiv:1112.4173v2 (2011).